

Post Treatment Instructions – Med Spa, Liquid Lift

Maximize your cosmetic results with a liquid lift treatment (ex. Botox, Dysport, Xeomin, hyaluronic acid, Restylane, Juvederm, Belotero, collagen, Radiesse, Sculptra, or injectable filler) by following these basic post-treatment instructions. Please contact the office with any questions.

General

- Numbness, bruising, slight redness, swelling, tenderness, or itching sensation around the treated area is common
- No smoking or alcohol, at minimum 24 hours
- No aspirin, ibuprofen, Motrin, Advil, or similar anti-inflammatory medication. You will be advised you when you may resume taking these medications after the procedure.
 - Other blood thinners, such as Coumadin or Plavix, must also be discontinued, under the guidance of your primary care physician.
- No herbal medications, supplements, or teas.
 - Increased risk of bleeding include, but are not limited to Vitamin E, garlic, ginger, ginkgo, ginseng, kava, and St. John's Wort, fish oil, and green tea
 - Arnica montana herbal *tablet* may help reduce bruising and swelling

Diet

- No restrictions on specific type of food or drink, *except* for lip injections then avoid hot food/drink. Drink plenty of fluids. Do not chew gum. You may use a straw.

Activity

- Sleep with your head elevated for the first 48 hours, to help reduce facial swelling.
- Do NOT blow nose.
- Avoid sneezing. If unable to avoid sneezing, then sneeze with your mouth open.
- Do NOT bend over or hang your head down.
- No heavy lifting, straining, or strenuous activity for at least 48 hours.
- Avoid any other facial treatments, including waxing and microdermabrasion for 48 hours.
- Light kissing is permitted.

Treatment Area Care

- Ice over treated area helps reduce facial swelling.
- Do not rub the area.
- Gently pat the area dry with a clean towel. Do not rub.
- Avoid direct sun exposure. You may resume sunscreen over the area immediately.
- Make-up may generally be used the same day following the procedure.
- *Sculptra only*: Deep facial massage area for 5 minutes, 5 times a day, for 5 days to stimulate collagen production.

Please notify Dr. Chaboki if you suffer from cold sores, as cold sores may occur with treatments around the mouth or lips.