

Postoperative Instructions – Cheek Augmentation, Chin Implant

Maximize your results after facial sculpting surgery with cheek augmentation or chin implant by following these basic post-treatment instructions. Please contact Dr. Chaboki with any questions.

General

- Numbness in the cheeks, lips, nose, and teeth is common
- No smoking or alcohol
- No aspirin, ibuprofen, Motrin, Advil, or similar anti-inflammatory medication. Dr. Chaboki will advise you when you may resume taking these medications.
 - Other blood thinners, such as Coumadin or Plavix, must also be discontinued, under the guidance of your primary care physician.
- No herbal medications, supplements, or teas.
 - Increased risk of bleeding include, but are not limited to Vitamin E, garlic, ginger, ginkgo, ginseng, kava, and St. John's Wort, fish oil, and green tea
 - Arnica montana herbal *tablet* may help reduce bruising and swelling.

Diet

- Advance slowly from liquids to very soft foods. No foods are allowed which require chewing for 1 week after surgery.
- Drink plenty of fluids. Do not chew gum. You may use a straw.

Activity

- Limit mouth or jaw opening and stretching for 1 week.
- Sleep with your head elevated for the first week, to help reduce facial swelling.
- Do NOT rub your nose, cheek, or chin, or put any pressure on these areas.
- Do NOT blow nose.
- Avoid sneezing. If unable to avoid sneezing, then sneeze with your mouth open.
- Do NOT bend over or hang your head down.
- No heavy lifting, straining, strenuous activity, or sex for at least 2 weeks.
- No contact sports for 6 weeks.

Hygiene

- Be very gentle with brushing your teeth for at least 2 weeks.
- Keep the treated areas dry when bathing.

Cheek

- Rinse mouth after every meal with 50-50 mix of water-hydrogen peroxide or your prescription mouth wash. Internal sutures will dissolve after a short period of time.
- Cotton bolsters and taping may be over the cheeks and will removed after 1-2 days.

Chin

- Remove your tape dressing after 2 days. Small amount of blood is normal.
- Place plain petrolatum ointment (ex. Vaseline) via a cotton-tip applicator on the incisions. Keep moist with ointment at all times for 1 month. *Do not use any other creams/lotions.*

Houtan Chaboki, MD
Board Certified Facial Plastic Surgeon
www.potomacplasticsurgery.com



2021 K Street, NW, Suite 206
Washington, DC 20006
202.741.3409

Specializing in rhinoplasty (nose job surgery), face lift, neck lift, blepharoplasty (eyelid surgery), brow lift, Botox, facial fillers, and skin care for Washington DC, Maryland, and northern Virginia.