

Postoperative Instructions – Face Lift and Neck Lift

Maximize your cosmetic results after facelift or neck lift surgery by following these basic post-treatment instructions. Please contact Dr. Chaboki with any questions.

General

- Numbness or stiffness around the neck, cheek, face, and ears is common
- Keep hair dryer on cool setting. Be very careful while brushing hair.
- Make-up may be used one week following the procedure
- No smoking or alcohol
- No aspirin, ibuprofen, Motrin, Advil, or similar anti-inflammatory medication. Dr. Chaboki will advise you when you may resume taking these medications.
 - Other blood thinners, such as Coumadin or Plavix, must also be discontinued, under the guidance of your primary care physician.
- No herbal medications, supplements, or teas.
 - Increased risk of bleeding include, but are not limited to Vitamin E, garlic, ginger, ginkgo, ginseng, kava, and St. John's Wort, fish oil, and green tea
 - Arnica montana herbal *tablet* may help reduce bruising and swelling

Diet

- Advance slowly from liquids to soft, then solid foods. No restrictions on specific type of food or drink. Drink plenty of fluids. Do not chew gum.

Activity

- Sleep with your head elevated for the first 48 hours, to help reduce facial swelling.
- Do NOT blow nose.
- Avoid sneezing. If unable to avoid sneezing, then sneeze with your mouth open.
- Do NOT bend over or hang your head down.
- No heavy lifting, straining, strenuous activity, or sex for at least 2 weeks.

Wound Care

- Remove your gauze dressing after 2 days. Small amounts of blood is normal.
- Keep clean ice-packs, wrapped in a clean cloth, over the area for the first 48 hours. Use ice every hour while awake to help reduce swelling.
- Steri-strip tape bandages may have been applied to your incision. Leave this tape in-place. The tape will gradually peel off in a few days.
- Place plain petrolatum ointment (ex. Vaseline) via a cotton-tip applicator on the incisions. Keep moist with ointment at all times for 1 month. *Do not use any other creams/lotions.*
- Keep the incision dry. You may wash the area with a mild soap and luke warm water *after 2 days*. Do not soak the area in water, or leave wet dressing against wound.
- Gently pat the area dry with a clean towel. Do not rub.
- Avoid direct sun exposure. Generally, you may start using sunscreen over the area at 1 week after the sutures are removed.
- Wear elastic neck support dressing continuously, removing only to apply ointment. Elastic neck support should *not* be tight, but rather provide a gentle “hug” to area.

Houtan Chaboki, MD
Board Certified Facial Plastic Surgeon
www.potomacplasticsurgery.com



2021 K Street, NW, Suite 206
Washington, DC 20006
202.741.3409

Specializing in rhinoplasty (nose job surgery), face lift, neck lift, blepharoplasty (eyelid surgery), brow lift, Botox, facial fillers, and skin care for Washington DC, Maryland, and northern Virginia.